

Characterization and Antimicrobial Susceptibility Patterns of Isolates from Ward Fomites

Farouk Segujja^{1,2*}, Bashir Mwambi², Charles Kato Drago¹,
Nathan Musisi Lubowa¹, Joseph Mugambwa³ and Patrick Wabuyi²

¹Department of Biomolecular Resources, College of Veterinary Medicine, Animal Resources and Biosecurity, Makerere University, P.O.Box 7062, Kampala, Uganda.

²Department of Medical Microbiology, International Health Sciences University, P.O.Box 7782, Kampala, Uganda.

³Department of Medical Microbiology, Faculty of Health Sciences, Habib Medical School, Islamic University in Uganda, Kampala Campus, P.O.Box 7689, Kampala, Uganda.

Authors' contributions

This work was carried out in collaboration between all authors. Author FS designed the study, wrote the protocol, participated in the data collection and wrote the first draft of the manuscript. Author BM performed the data entry and statistical analysis. Authors CKD and NML supervised and conceptualized the project. Authors JM and PW managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/BBJ/2016/27861

Editor(s):

(1)

(2)

Reviewers:

(1)

(2)

Complete Peer review History:

Original Research Article

Received 22nd June 2016

Accepted 10th July 2016

Published 20YY

ABSTRACT

Aim: The study was conducted to determine antimicrobial susceptibility patterns among isolates from ward fomites at Kiwoko Hospital and to detect resistances in the form of Macrolide Lincosamide StreptograminB (MLS_B), Methicillin Resistant *Staphylococcus aureus* (MRSA), Extended Spectrum β Lactamases (ESBLs), *AmpC*, and Multi Drug Resistant (MDR) pathogens.

Study Design: Laboratory based cross-sectional study.

Place and Duration of Study: The study was conducted in various wards and sections at Kiwoko Hospital, a rural setting in the central region of Uganda, between January and June 2015.

Methodology: We recruited 290 samples from the Surgical, Medical, Maternity and Pediatric

*Corresponding author: E-mail: segujjaf@gmail.com;

wards as well as the Out Patient Department (OPD) at Kiwoko Hospital for the study. Samples were taken by swabbing the different surfaces and instruments which included; sphygmomanometers, stethoscopes, beds, nurses' stations, staff/visitors' chairs, door handles, patients' crepe bandages, curtains, switches, and sink handles among others. Susceptibility testing was done using the disc diffusion methods by Kirby Bauer for phenotypic expression of MLS_B resistances, MRSA, MSSA, ESBL, MDR and AmpC. Co-resistances exhibited by isolated ESBL producers were also phenotypically tested.

Results: Of the 290 surfaces and instruments swabbed, 57.59% (CI= 49.18 - 67.01) carried bacterial pathogens and by using standard surface agar plating methods, *Staphylococcus aureus* was the mostly isolated pathogen 43 (25.75%), followed by *Klebsiella pneumoniae* 35 (20.96%), *Escherichia coli* 31 (18.55%), *Pseudomonas aeruginosa* 20 (11.98%), *Enterococcus faecalis* 12 (7.19%), *Staphylococcus epidermidis* 10 (5.98%), *Proteus mirabilis* 9 (5.39%), *Bacillus* spp. 4 (2.40%), and *Staphylococcus saprophyticus* 3 (1.80%). Among enterobacteriaceae, 5 (6.67%, CI= 2.16 – 15.56) were identified as AmpC producers and 16 (21.33%, CI= 12.19 - 34.64) as ESBL producers out of which 4/16 (25.00%, CI = 6.81 – 64.01) showed ESBL co-resistance. Of the 43 *Staphylococcus aureus* isolates, 9.30% were MRSA (CI = 2.53 - 23.82) and 90.70% MSSA (CI = 64.49 - 100). In MLS_B resistance patterns, 23.26% of the total *S. aureus* isolates were constitutive MLS_B while 6.98% showed inducible MLS_B as 27.91% exhibited an MS phenotype. Out of all the isolates recovered from fomites, 27/167 (16.17%, CI = 10.65 – 23.52) were identified as Multi Drug Resistant (MDR).

Conclusion: Hospital fomites harbored resistant pathogens that could well persist for a long period of time thereby predisposing patients to Hospital acquired infections. Therefore, routine screening of clinical samples for MLS_B, ESBL, AmpC, MRSA and MDR could significantly monitor potential treatment failures in the management of resistant bacterial infections spread by pathogens on ward items and surfaces at Kiwoko Hospital, Uganda.

Keywords: Nosocomial infections; Macrolide Lincosamide StreptograminB; phenotypic; ward; fomites.

ABBREVIATIONS

ATCC	: American Type Culture Collection.
CDH	: Central Drug House.
CLSI	: Clinical and Laboratory Standard Institute.
cMLS _B	: Constitutive Macrolide-Lincosamide-StreptograminB.
ESBL	: Extended Spectrum β Lactamase.
HAIs	: Hospital Acquired Infections.
iMLS _B	: Inducible Macrolide Lincosamide-StreptograminB.
MDR	: Multi-Drug Resistant .
MRSA	: Methicillin-Resistant <i>Staphylococcus aureus</i> .
MS	: Macrolide-Streptogramin B resistant phenotype.
MSSA	: Methicillin-Sensitive <i>Staphylococcus aureus</i> .
AMPC	: This is a chromosomal mediated, inducible mechanism of resistance to Cephamycins and 3rd generation spectrum as a result of acquisition of AMPC gene or depression of a chromosomal AMPC gene.
AMP C Inducer	: An antimicrobial that switches on the AMPC resistance gene to render treatment with cephamycins and 3 rd generation cephalosporin inappropriate.
ESBL	: Extended spectrum β-lactamase (ESBL) is an enzyme whose coding gene is a bacterial Plasmid and confers resistance to β-lactam antibiotics up to the 3 rd generation cephalosporins, penicillins and aztreonams thereby rendering them inactive.
FOMITES	: Objects or materials which are likely to carry or harbour pathogens that cause infections in a healthcare setting/hospital.
MLS resistance	: Resistance to macrolides such as erythromycin, and lincosamides such as clindamycin, usually is due to an erm gene. These erm genes code for production of an RNA methylase enzyme that modifies the ribosomal binding site of macrolides, lincosamides, and StreptograminB.

1. INTRODUCTION

Although its clearly evident that several control measures including use of checklists, effective handwashing, surveillance and setting up infection control units have been adopted in many healthcare settings over the years to help reduce rates of contamination [1,2], hospital acquired infections also known as nosocomial infections remain a major cause of increased patient management costs [3,4], prolonged hospital stays and death among hospitalized patients worldwide [5-8].

Nosocomial infections are infections occurring in hospitalized patients in whom the infection was not present or incubating at the time of admission, these encompass infections acquired in the hospital but appearing after discharge, and also occupational infections among staff of the facility [7]. At any one moment, approximately 5% to 10% of hospitalized patients have an infection acquired after admission [9]. Higher nosocomial infection rates have been reported in developing countries more often than those documented from developed countries [4,6,7,10,11].

Recent studies suggest that contaminated environmental surfaces may play a major role in the transmission of nosocomial infections [12-15]. Harbored organisms may well survive for long periods on items or surfaces thereby presenting as a major health concern [5,6,10,16,17]. Such inanimate surfaces or objects are often known as fomites and are greatly associated with detrimental infection outbreaks [18-20].

Despite undoubtable proof that fomites do harbor nosocomial causing pathogens [5,12,13,16], there still remains substantial controversy on whether or not some healthcare personnel contribute to the transmission of these infections in hospital settings. Several studies have however indicated that healthcare workers are equally involved and that bacterial contamination of their hands is likely to result into transmission to patients [14,21-23]. Without contact, pathogens can still find their way to a new host via contaminated air by patients carrying airborne infections commonly through coughing and sneezing [24].

In an attempt to treat nosocomial infections, there has been increased use of

first-line antibiotics where appropriate second-line drugs are not readily available or affordable, this has eventually led to the emergence of quite a number of resistant bacteria that have subsequently persisted in the hospital environment thereby becoming endemic [7]. Many resistant strains including ESBL producers [25], AmpC producers [26,27], multidrug resistant pathogens [28,29], *Pseudomonas aeruginosa* [30], *Enterococcus* spp. [5,16,17] and MLS_B resistant isolates [31-33], have been prevalently reported in many hospital settings as potential life threatening pathogens worldwide.

Little is known about bacterial contamination of the environment around patients admitted to local hospitals in Uganda. This study therefore aimed at investigating the presence, distribution, rate and antibiogram of pathogens isolated from ward items and surfaces of Kiwoko Hospital that could potentially predispose patients to hospital acquired infections.

2. MATERIALS AND METHODS

2.1 Study Design

This was a laboratory based descriptive cross sectional study conducted between January and June of 2015 at Kiwoko Hospital and Medical Microbiology laboratories of Makerere University, International Health Sciences University and Habib Medical School, Islamic University In Uganda, Kampala Campus.

2.2 Study Samples

These were swabs taken from all items and surfaces in surgical, medical, pediatric and maternity wards as well as OPD sections at Kiwoko Hospital. Disinfection was always done at the start of the day and samples were collected during work. Fomites included; thermometers, sphygmomanometer, stethoscopes, visitors' chairs, intravenous poles, beds, bedside tables, hand towels, kidney dishes, patients' crepe bandages, curtains, telephones, dressing trolleys, light switches, door handles, record books, recording pens, examination bed area, pediatric weighing scale, pulsometer control panels, cupboard shelves, drug locks, screens, cardiocograph control panels, counter tops (Nurses and doctors' stations) and water sinks (Patients' sinks, Doctors' sinks, Nurses' sinks).

2.3 Laboratory Procedures

2.3.1 Specimen collection and identification of pathogen

Specimen collection involved using a sterile cotton swab stick moistened with 0.9% physiological saline. The swab was then pressed and rolled several times over the entire surface of a selected item. With care not to touch the swab, it was then immediately placed into an appropriate bijoux bottle containing Brain heart infusion broth and aerobically incubated overnight at 37°C as published by Bauer et al [34].

On suspected growth exhibited by turbidity in the broth, specimens were subcultured on Blood agar (Central Drug House (CDH), India) and MacConkey agar (CDH, India) using aseptic streaking techniques followed by 24 hours of incubation at 37°C aerobically. Plates were then read for bacterial growth, and organisms examined for their characteristic colonial appearance, hemolysis, swarming, and/or pigmentation on the different media, before subsequent follow-up for identification and confirmation through gram-staining, sugar fermentation, and biochemical reactions. On failure to grow within 24 hours, plates were further re-incubated for the same time under similar conditions before discarding them and recording their results as negative.

Members of the family Enterobacteriaceae were identified by indole production, Hydrogen Sulphide (H₂S) production, citrate utilization, gas production, motility tests, urease test, oxidase, and carbohydrate utilization. For gram-positive bacteria identification and confirmation, coagulase, DNase, catalase, mannitol fermentation, CAMP (Christie, Atkins, and Munch Peterson) test, esculin bile test, bacitracin, optochin and novobiocin susceptibility tests were all used.

2.3.2 Antimicrobial susceptibility testing

Antimicrobial susceptibilities were studied by modified Kirby-Bauer's agar disc diffusion methods [34] and according to guidelines from Clinical and Laboratory Standards Institute [35]. Susceptibility testing was done in categories for Enterobacteriaceae (*Klebsiella pneumoniae*, *E. coli* and *P. mirabilis*), non-enterobacteriaceae (*P. aeruginosa*), *S. aureus* and *E. faecalis*.

Testing was not done for isolated *S. epidermidis*, *S. saprophyticus*, and *Bacillus* spp.

Among the enterobacteriaceae isolates, antibiotics tested included; Augmentin (AUG, 30 µg), Ceftriaxone (CRO-30 µg), Cefoxitin (FOX, 30 µg), Ertapenem (ETP, 10 µg), Imipenem (IPM, 10 µg) and Piptazocin (PTZ, 110 µg) a drug combination of piperacillin and tazobactam.

These six discs phenotypically identified ESBL and *AmpC* producing enterobacteriaceae by positioning the Augmentin disc in the centre of the media plate at a distance of 20 mm away from a 3rd generation cephalosporin (Ceftriaxone, 30 µg) on one side, and Cefoxitin (acting as a strong labile *AmpC* inducer) on the other (double synergism method). As a last resort treatment option for ESBLs and AMPC pathogens, imipenem (IPM, 10 µ) was included to ascertain whether or not the isolate was Carbapenem Resistant Enterobacteriaceae (CRE), this imipenem also acted as another AMPC inducer. To report CRE results correctly, another stronger carbapenem drug in the form of Ertapenem (ETP, 10 µg) was included as per the CLSI defining guidelines [35].

Potential ESBL producers showed synergism between clavulanic acid (AUG, 30µg) and ceftriaxone (3rd Generation cephalosporin), resistance to Piptazocin (PTZ, 110µg) and sensitivity to Cefoxitin (FOX, 30µg), a cephamycin [26,35,36].

AmpC gene producing enterobacteriaceae exhibited resistance to all Augmentin (AUG, 30 µg), Ceftriaxone (CRO, 30 µg), Cefoxitin-30 µg, and also showed a flattening (blunting) of the zone size of Ceftriaxone at its junction with the zone edges of both Cefoxitin and Augmentin that were adjacently placed 20 mm away from it on either sides [37].

Sensitivity testing was further done on pathogens that showed ESBL and *AmpC* production to determine which drugs would be used as treatment options and to ascertain whether they still showed co-resistance patterns against such alternatives. These drug categories included; an aminoglycoside (Gentamycin, 10 µg), Fluoroquinolones (Ciprofloxacin, 10 µg), Chloramphenicol (Chloramphenicol, 30 µg), Folic acid inhibitor (Sulphamethazole-Trimethoprim, 1.25 µg), or Tetracyclines (Tetracycline, 30 µg) [25,36]. An ESBL producer that showed

resistance to at least one agent in three or more antimicrobial categories for which it did not have known intrinsic resistance was defined as a co-resistant pathogen [29].

Among the non-enterobacteriaceae (particularly *P. aeruginosa*), antimicrobial agents tested included; Ceftazidime (CAZ, 30 µg), Meropenem (MEM, 10 µg), Gentamycin (CN, 10 µg), Ciprofloxacin (CIP, 30 µg) and Piptazocin (PTZ, 110) a drug combination of piperacillin and tazobactam. These were kept five to limit synergism that would falsely show increased zone size diameters between Piptazocin and ciprofloxacin, or piptazocin and Gentamycin [38].

For *Enterococcus faecalis*, antibiotic discs tested were; Gentamycin (CN, 10 µg), Vancomycin (VA, 30 µg) to detect Vancomycin Resistant *Enterococcus* (VRE), Clindamycin (DA, 2 µg), Erythromycin (E, 15 µg), Ceftriaxone (CRO, 30 µg), and Ciprofloxacin (CIP, 30 µg).

In testing susceptibility patterns of *S. aureus*, six (6) discs were set including Cefoxitin - FOX, 30 µg (as a stronger surrogate to oxacillin or methicillin), Gentamycin (CN, 10 µg), Vancomycin (VA, 30 µg) to detect Vancomycin Resistant *S. aureus* (VRS) and as a treatment option for MRSA, Ciprofloxacin (CIP, 10 µg), Erythromycin (E, 15 µg) and Clindamycin (DA, 2 µg).

To detect Macrolide-Lincosamide-StreptograminB (MLS_B) resistances, Erythromycin and Clindamycin discs were placed at a distance of 15mm edge to edge from each other [39]. A positive "D test" showed flattening of the zone of inhibition of Clindamycin (D-shaped inhibition zone ≥21 mm) at the area besides which it was adjacent to Erythromycin (zone size ≤ 13 mm) and was defined as an inducible MLS_B (iMLS_B) phenotype. *S. aureus* strains that were resistant to both Erythromycin and Clindamycin with a perfect circular-like inhibition zones were defined as constitutive MLS_B (cMLS_B) phenotype. Isolates that were resistant to Erythromycin (zone size ≤13 mm) but sensitive to Clindamycin (zone size ≥21 mm) without D-shaped zones around Clindamycin were defined as MS phenotypes [32].

A multidrug resistant (MDR) phenotype of isolates was identified as an expression of resistance to at least one agent in three or more different antimicrobial categories to which these isolates did not have known intrinsic resistances.

These MDR defining categories included: cephamycin, cephalosporins, carbapenems, glycopeptides, aminoglycosides, macrolides, lincosamides, fluoroquinolones, folate pathway inhibitors (sulfamethazole-trimethoprim), antipseudomonal Penicillin-β-lactamase inhibitors (piptazocin), tetracycline, and chloramphenicol. All MRSA were defined as MDR by virtue of being MRSA as this predicted resistance to cephamycin (Cefoxitin, a surrogate marker for Methicillin), ciprofloxacin, and all β lactam antibiotics [29].

2.4 Quality Control

Reference strains used as controls were: *E. coli* (ATCC 25922), *E. faecalis* (ATCC 29212), *S. aureus* (ATCC 25923), Methicillin Resistant *S. aureus* (ATCC 43300), *Escherichia coli* ATCC 35218 (ESBL producer), *K. pneumoniae* ATCC 700603 (ESBL producer), and *P. aeruginosa* (ATCC 27853).

2.5 Data Analysis

Data was entered in Microsoft excel, cleaned and imported to Stata version 13 statistical package for analysis. Results were then presented in form of graphs and tables for proportions, absolute values, percentages, and confidence intervals calculated by Poisson's test for point estimates at 95% level of confidence with a *P*-value of 0.05 considered as statistically significant.

3. RESULTS AND DISCUSSION

During the study period, 290 items and surfaces from across all wards were recruited. Among these, 167 (57.59%, CI = 49.18 - 67.01) harbored bacterial pathogens. *S. aureus* (25.75%) and *K. pneumoniae* (20.96%) were the most frequently isolated bacteria, followed by *E. coli* (18.55%), *P. aeruginosa* (11.98%), *E. faecalis* (7.19%), *S. epidermidis* (5.98%), *P. mirabilis* (5.39%), *Bacillus* species (2.40%) and *S. saprophyticus* (1.80%) (Table 1).

3.1 Antimicrobial Susceptibility Profiles of Isolates from Ward Fomites

3.1.1 Antibiogram of enterobacteriaceae isolates

Amongst Enterobacteriaceae, different isolates showed varying susceptibility patterns to the set of first antibiotics that included: Augmentin (AUG, 30 µg), Ceftriaxone (CRO, 30 µg), Imipenem

(IPM, 10 µg), Ertapenem (ETP, 10 µg), Cefoxitin (FOX, 30 µg), and Piptazocin (PTZ, 110 µg). The most effective antibiotics against enterobacteriaceae (*P. mirabilis*, *E. coli* and *K. pneumoniae*) were imipenem and Ertapenem with a sensitivity rate of 100% (Table 2).

3.1.1.1 Phenotypic detection of AmpC and ESBL among enterobacteriaceae

With susceptibility patterns exhibited by enterobacteriaceae (Table 2), ESBL and AmpC producers were phenotypically detected (Table 3). ESBL producing enterobacteriaceae showed synergism between clavulanic acid (AUG, 30 µg) and ceftriaxone (3rd Generation

cephalosporin). They also expressed resistance to Piptazocin (PTZ, 110 µg) and sensitivity to Cefoxitin (FOX, 30 µg) a cephamycin.

AmpC gene producing enterobacteriaceae exhibited resistance to Augmentin (AUG, 30 µg), Cefoxitin-30 µg.

Out of the 75 enterobacteriaceae isolates, 16 (21.33%, CI= 12.19 - 34.64) were identified as ESBL producers while 5 (6.67%, CI= 2.16 - 15.56) phenotypically emerged as AmpC producers. *K. pneumoniae* and *P. mirabilis* expressed the highest resistance genes (ESBL and AmpC) amongst all enterobacteriaceae (Table 3).

Table 1. Bacterial profile of ward fomites at Kiwoko Hospital

Isolate	Frequency	Percentage	95% CI
<i>S. saprophyticus</i>	3/167	1.80%	0.37 - 5.25
<i>Bacillus</i> species	4/167	2.40%	0.65 - 6.13
<i>P. mirabilis</i>	9/167	5.39%	2.46 - 10.42
<i>S. epidermidis</i>	10/167	5.98%	2.87 - 11.01
<i>E. faecalis</i>	12/167	7.19%	3.71 - 12.55
<i>P. aeruginosa</i>	20/167	11.98%	7.32 - 18.50
<i>E. coli</i>	31/167	18.55%	12.61 - 26.35
<i>K. pneumoniae</i>	35/167	20.96%	14.60 - 29.15
<i>S. aureus</i>	43/167	25.75%	18.63 - 34.68
Total	167/290	(57.59%)	49.18 - 67.01

*Bacterial pathogen recovered from ward items, CI - Confidence Interval

Table 2. Antibiogram of isolated enterobacteriaceae

Antibiotics	Isolates							
	<i>Escherichia coli</i> n= 31		<i>Proteus mirabilis</i> n= 9		<i>Klebsiella pneumoniae</i> n= 35		Total n= 75	
	R (%)	S (%)	R (%)	S (%)	R (%)	S (%)	R (%)	S (%)
Cefoxitin	3 (9.68)	28 (90.32)	1 (11.11)	8 (88.89)	5 (14.29)	30 (85.71)	9 (12)	66 (88)
Augmentin	13 (41.94)	18 (58.06)	5 (55.56)	4 (44.44)	16 (45.71)	19 (54.29)	34 (45.33)	41(54.67)
Imipenem	0 (0.00)	31 (100)	0 (0.00)	9 (100)	0 (0.00)	35 (35)	0 (0.00)	75 (100)
Ertapenem	0 (0.00)	31 (100)	0 (0.00)	9 (100)	0 (0.00)	35 (100)	0 (0.00)	75 (100)
Ceftriaxone	13 (41.94)	18 (51.43)	4 (44.44)	5 (55.56)	18 (58.06)	17 (48.57)	35 (46.67)	40(55.33)
Piptazocin	14 (45.16)	17 (54.84)	4 (44.44)	5 (55.56)	15 (42.86)	20 (57.14)	33 (44)	42 (56)
Total (%)	43 (23.12)	143(76.88)	14(25.93)	40(74.07)	54 (25.71)	156(74.29)	(24.67)	(75.33)
CI (%) =	16.73-31.14	64.80-90.57	14.17-43.5	52.92-100	19.32-3.55	63.06- 86.90	20.29-29.7	67.53-83.8

*R – Resistant, S – Sensitive, CI – Confidence Interval

Table 3. Prevalence of ESBL and AmpC producing enterobacteriaceae

Resistance definition	Isolates			Total Enterobacteriaceae n = 75, (%)
	<i>Klebsiella pneumoniae</i> n = 35	<i>Escherichia coli</i> n = 31	<i>Proteus mirabilis</i> n = 9	
ESBL producer	8 (25.81%)	5 (16.13%)	3 (33.33%)	16(21.33%)CI=12.19-34.64
AmpC producer	3 (8.57%)	2 (6.45%)	0 (0.00%)	5 (6.67%) CI= 2.16 -15.56
Total (%)	11 (31.43)	7 (22.58)	3 (33.33)	21 (28.0)
CI (%) =	CI=15.69 – 56.23	CI=9.08– 46.52	CI=6.87-97.41	CI= 17.33 - 42.80

ESBL – Extended spectrum β lactamase, CI – Confidence Interval

3.1.1.2 Co-resistance among ESBL enterobacteriaceae

Out of the 16 ESBL producing enterobacteriaceae isolated, 4 (25%, CI= 6.81 - 64.01) showed co-resistance. Among these, 2 were identified as *K. pneumoniae* 2(12.5%, CI= 1.51- 45.15) was the most frequently isolated co-resistant pathogen followed by *E. coli* (6.25%) and *P. mirabilis* (6.25%) (Fig. 1).

The most effective drugs for ESBL producing enterobacteriaceae were carbapenems (imipenem and ertapenem) as shown in Table 2, followed by Ciprofloxacin at a low resistance rate of 6.25% (CI= 0.16 – 34.82), Gentamycin (12.5%) and Chloramphenicol (12.5%). The most antimicrobial resistances in ESBL producers were reported with Tetracycline (37.5%, CI= 13.76 - 81.6) and Sulphamethazole-Trimethoprim (31.25%, CI= 10.15 - 72.93) (Table 4).

3.1.2 Antibiogram of non enterobacteriaceae isolates

The susceptibility rate of *P. aeruginosa* was highest for Piptazocin (100%), a drug combination of piperacillin and tazobactam. It

showed utmost resistance to ciprofloxacin (90%) (Table 5).

3.1.3 Antibiogram of Enterococcus faecalis

E. faecalis showed 100% resistance to 50% of the set antibiotics leaving Vancomycin and Gentamycin as the only treatment options at sensitivity levels of 100% and 75% respectively (Table 6).

3.1.4 Antimicrobial susceptibility pattern of Staphylococcus aureus

The susceptibility rate of all *S. aureus* isolates was highest for Vancomycin (100%) irrespective of whether they were identified as Methicillin resistant or Methicillin Sensitive *S. aureus* (Table 7).

3.1.4.1 Phenotypic detection of MLS_B resistances and MRSA

Out of all the 43 *S. aureus* pathogens isolated from ward fomites, 39 (90.70%) were detected as Methicillin Sensitive *S. aureus* (MSSA) while 4 (9.30%, CI= 2.53 - 23.82) were Methicillin Resistant *S. aureus* (Table 8).

Table 4. Antibiogram showing treatment options for ESBL enterobacteriaceae

Antibiotics	ESBL producing enterobacteriaceae (n=16)					
	<i>Escherichia coli</i> n=5		<i>Proteus mirabilis</i> n=3		<i>Klebsiella pneumoniae</i> n=8	
	R (%)	S (%)	R (%)	S (%)	R (%)	S (%)
Tetracycline	1(20.00)	4(80.00)	3(100.0)	0(0.00)	2(25.00)	6(75.00)
Gentamycin	1(20.00)	4(80.00)	0(0.00)	3(100.0)	1(12.50)	7(87.50)
Chloramphenicol	0(0.00)	5(100.00)	1(33.33)	2(66.67)	1(12.50)	7(87.50)
Ciprofloxacin	0(0.00)	5(100.00)	0(0.00)	3(100)	1(12.50)	7(87.50)
Sulpha-trimethoprim	1(20.00)	4(80.00)	1(33.33)	2(66.67)	3(37.50)	5(62.50)
Total (%)	3(12.00)	22(88.00)	5(33.33)	10(66.67)	8(20.00)	32(80.00)
CI (%) =	2.47-35.07	55.15-100	10.82-7.79	31.97-100	8.63-39.41	54.72-100

R – Resistant S – Sensitive CI – Confidence Interval, ESBL – Extended spectrum β lactamase, CI – Confidence Interval




Fig. 1. Co-resistance among ESBL enterobacteriaceae
ESBL – Extended spectrum β lactamase

Table 5. Susceptibility pattern of isolated non-enterobacteriaceae

Antibiotics	Isolate	
	<i>Pseudomonas aeruginosa</i> n = 20	
	R (%)	S (%)
Ceftazidime (CAZ, 30 µg)	6 (30.00)	14 (70.00)
Meropenem (MEM, 10 µg)	7 (35.00)	13 (65.00)
Gentamycin (CN, 10 µg)	14 (70.00)	6 (30.00)
Ciprofloxacin (CIP, 30 µg)	18 (90.00)	2 (10.00)
Piptazocin (PTZ, 110 µg)	0 (0.00)	20 (100.00)
Total (%)	45 (45.00)	55 (55.00)
CI (%) =	32.82 - 60.21	41.43 - 71.60

R – Resistant S – Sensitive CI – Confidence Interval

Out of the 25 (58.14%, CI= 37.62 – 85.83) erythromycin resistant *S. aureus* isolates, 3 (12%, CI= 2.47 – 35.07) expressed an inducible MLS_B phenotypic resistance pattern in which D-shaped inhibition zones of Clindamycin placed 15mm away from erythromycin were observed. A total of 10 (40%, CI= 19.18 – 73.56) *S. aureus* isolates were identified as constitutive MLS_B with resistances to both erythromycin and clindamycin. All the 4 isolated MRSA exhibited a constitutive MLS_B pattern while 12 (48%, CI= 24.86 – 83.85) isolates presented with an MS resistant phenotype (Table 8).

Table 6. Antimicrobial susceptibility profile of *Enterococcus faecalis*

Antibiotics	Isolate	
	<i>Enterococcus faecalis</i> n = 12	
	R (%)	S (%)
Vancomycin (30 µg)	0 (0.00)	12 (100.00)
Clindamycin (2 µg)	12 (100.00)	0 (0.00)
Erythromycin (15 µg)	12 (100.00)	0 (0.00)
Gentamycin (10 µg)	3 (25.00)	9 (75.00)
Ceftriaxone (30 µg)	12 (100.00)	0 (0.00)
Ciprofloxacin (30 µg)	5 (41.67)	7 (58.33)
Total (%)	44 (61.11)	28 (38.89)
CI (%) =	44.40 - 82.04	25.84 - 56.21

R – Resistant S – Sensitive CI – Confidence Interval

Table 7. Antibigram of *Staphylococcus aureus* isolated from hospital fomites

Antibiotic	Isolate	
	<i>Staphylococcus aureus</i> n = 43	
	R (%)	S (%)
Cefoxitin (30 µg)	4(9.30)	39(90.70)
Clindamycin (2 µg)	13(30.23)	30(69.77)
Erythromycin (15 µg)	25(58.14)	18(41.86)
Gentamycin (10 µg)	18(41.86)	25(58.14)
Ciprofloxacin (30 µg)	12(27.91)	31(72.09)
Vancomycin (30 µg)	0(0.00)	43(100.00)
Total (%)	72 (27.91)	186 (72.10)
CI (%) =	21.84 - 35.14	62.10 - 83.23

R – Resistant S – Sensitive CI – Confidence Interval

Table 8. Distribution of MRSA and MLS_B resistances among *S. aureus*

Resistance definition	MSSA (%)	MRSA (%)	Total (%)	Antibiotic agent	
				E, 15 µg	DA, 2 µg
cMLS _B	6 (15.38)	4 (100.00)	10 (23.26)	R	R
iMLS _B	3 (7.69)	0 (0.00)	3 (6.98)	R	S
MS phenotype	12 (30.77)	0 (0.00)	12 (27.91)	R	S
No DA resistance	18 (46.15)	0 (0.00)	18 (41.86)	S	S
Total (%)	39 (90.70)	4 (9.30)	43		
CI (%) =	64.49 – 100	2.53 - 23.82			

R- Resistant, S- Sensitive, E- Erythromycin, DA- Clindamycin, MRSA - Methicillin Resistant *Staphylococcus aureus*
MSSA - Methicillin Sensitive *Staphylococcus aureus*, MLS_B - Macrolide Lincosamide Streptogramin B
cMLS_B - constitutive MLS_B; iMLS_B - inducible MLS_B, MS - Macrolide Streptogramin B; CI - Confidence interval

Table 9. Prevalence of multi-drug resistance among isolates

Isolate n = 167	Antimicrobial categorical resistance definition, MDR (%)	Confidence interval (%)
<i>P. aeruginosa</i> (n=20)	8 (40.00)	17.27 - 78.81
<i>K. pneumoniae</i> (n=35)	6(17.14)	6.29 - 37.31
<i>P. mirabilis</i> (n=9)	3(33.33)	6.87 - 97.41
<i>E. coli</i> (n=31)	4(12.90)	3.51 - 33.04
<i>S. aureus</i> (n=43)	6(13.95)	5.12 - 30.37
Total (%)	27/167 (16.17)	10.65 - 23.52

MDR - Multidrug resistant

3.1.5 Multi-drug resistance among isolates on ward fomites

The overall Multi-drug resistance rate was 16.17% (CI= 10.65 – 23.52) with *P. aeruginosa* singly accounting for 29.63% (8/27, CI= 12.79 – 58.38) of the total MDR prevalence.

3.2 Discussion

Our study found an overall bacterial contamination rate of 57.59% (167/290, 95%CI= 49.18 - 67.01%) that was obtained from all the fomites swabbed. The contamination of fomites in this study could be attributed to irregular disinfection, the types of disinfectants used, hygienic conditions, overcrowding and lack of surveillance and implementation of HAI control protocols [4,10]. Elsewhere, contamination has been documented to occur on fomites but in varying degrees. The contamination rate from our study was similar to the 44% Obtained at Cleveland Veterans Affairs Medical Centre, USA [12]. This could be attributed to the similarities of sampled items.

When compared to other studies elsewhere, the prevalence rate in our study was higher than that reported at Murtala Mohammed Specialist Hospital in Nigeria 23.33% (95% CI= 20.28 - 26.71%) [40], and at Al Imam Ali Hospital in Bagdad 4% [41]. However, it was significantly lower than what was reported at San Vicente Foundation, a teaching hospital in Columbia (98.7%; 157/159) [13], and Gondar University Hospital in which 83.1% (95% CI = 68.78 – 99.52%) of inanimate objects were contaminated [42].

The items swabbed at Murtala Mohammed Specialist Hospital were from theatre [40] contrary to our study that involved samples from various general wards. There are always high safety precaution measures taken to control infection transmission from theater fomites compared to general wards [43,44]. In theaters, there is always limited access to only authorized health workers, strict handwashing and wearing of sterile theater attires (gowns, surgical masks, surgical boots and gloves) [21,43,44]. This is contrary to the practice in general wards. This could be the reason for the higher rates of contamination in our study compared to what Nwankwo and colleagues reported in 2012 [40]. On the other hand, the lower prevalence rate in our study compared to what was documented at San Vicente [13] could be attributed to the

difference in settings. San Vicente is a teaching hospital, up till now several authors have reported that bacterial contamination is more common in teaching hospitals than in non-teaching [4,10,41]. In teaching hospitals, students are normally being introduced to ward sections and know less about safety measures taken to control bacterial contamination [45,46].

The obtained fomite contamination rate highlighted a major role of ward items as potential vehicles and reservoirs of nosocomial infections, this accords with findings of several previous bacterial investigations of fomites [5,16, 21-23,47]. We never assessed for the relatedness of clinical isolates to fomites in this study, however, numerous authors have reported HAIs from various global regions [4,6,10, 20,48,49] with some documenting genotypic similarities of clinical isolates to hospital surfaces [50,51]. Therefore, bacterial contamination of fomites in our study reflects a regular daily risk of exposure to several hospital-acquired infections.

Staphylococcus aureus was the most frequently isolated pathogen across all wards, a result consistent with findings from different studies across the globe [15,52]. This is attributed to the fact that it is a normal flora of the skin [53-55] that subsequently gets into contact with ward surfaces from which it eventually invades patients thereby causing infections especially on surgical sites and other life-threatening diseases [55,56]. Previous studies have documented nasal carriage of *S. aureus* with or without any pathogenic events [42,57-59]. In health adults, the percentage carriage of *S. aureus* in nostrils is estimated to be 40% [53]. The presence of *Proteus*, *K. pneumoniae* and *E. coli* which are enteric bacteria [53,55], was indicative of fecal pollution and poor personal hygiene especially irregular handwashing practice. As a normal flora, *P. aeruginosa* has been documented to inhabit nonsterile areas on healthy individuals, however, it is able to infect any tissues especially in immune-compromised patients [54,56], its presence on various ward surfaces and instruments could therefore provoke severe infections in the form of wound contamination following surgical procedures, or those associated with catheterization such as urinary tract complications that may well persist as a result of bio-film formations in urogenital organs [30].

Vancomycin was the most effective antimicrobial agent against *S. aureus* followed by Cefoxitin,

Ciprofloxacin, Clindamycin, Gentamicin and Erythromycin with susceptibility rates of 100%, 90.70%, 72.09%, 69.77%, 58.14% and 41.86% respectively as seen in Table 7. This pattern is consistent with susceptibilities of clinical isolates elsewhere [58,60]. Several Clindamycin resistances were identified in our study (Table 8) a finding quite comparable with a number of previous investigations that have highlighted their existence in hospital settings, however, clindamycin that showed a moderate sensitivity level (69.77%) in our study, remains highly recommended in the treatment of various staphylococcal resistant strains today because of its better oral bioavailability, low cost, its importance as an alternative antibiotic in penicillin-allergic patients, excellent tissue penetration, and the fact that it accumulates in abscesses [33,39] than the Cefoxitin and Ciprofloxacin (which performed better in this study). Vancomycin was remarkably excellent, however, it should be given when therapeutic options are limited due to severe renal impairment associated with its prolonged administration [61,62].

Our study demonstrated presence of cMLS_B resistant phenotypes in *S. aureus*, and therefore should a patient pick a *Staphylococcus aureus* pathogen from the hospital, then chance was 30.23% (95% CI= 16.10 – 51.70) that it would be resistant (cMLS_B and iMLSB) to clindamycin a commonly used antibiotic in this setting. Of all the MLS_B phenotypic resistances exhibited by *S. aureus* (25/43), clindamycin would actually treat 48% (12/25). This accords with several other scientists who have continuously recommended that clindamycin should be cautiously used in the treatment of MLS_B resistances because of its increasing failures [39,63,64]. The fact that clindamycin is bacteriostatic, many studies have disapproved of its use in the treatment of endovascular infections like endocarditis or septic thrombophlebitis all pointing to prescription issues [63].

This study shows that out of the 43 isolates of *S. aureus*, 4 (9.03%) were MRSA and these were only sensitive to Vancomycin (100%). Such a finding is empirically suggestive of treatment with Vancomycin in case of suspected MRSA as previously recommended by other scientists [7,35,60,63,65]. However, recent reviews report increasing Vancomycin Resistant *Staphylococcus aureus* (VRSA) strains [66]. Therefore, this does not only necessitate the

need to seek other VRSA therapeutic options (Linezolid, daptomycin or ceftaroline), but also routine screening of clinical samples for staphylococcal antimicrobial resistance to these options as a better patient management strategy in rural hospitals.

E. faecalis showed the highest resistance rate of all isolates 61.11% (95% CI= 44.40 - 82.04%) as seen in Table 6. *E. faecalis* is known to be intrinsically resistant to a number of antimicrobial classes [29,67] and this therefore explains its exceptionally high insensitivity to the antimicrobial agents in this present study. Vancomycin was the most effective antibiotic against *E. faecalis* with a sensitivity rate of 100%, a finding similar to several earlier studies that have endorsed it as the best therapeutic drug of choice against *Enterococcus* species [7,35]. However, transferable resistance to vancomycin is now prevalently common in *Enterococcus* and has found its way into MRSA strains [66,68], thereby making treatment of Vancomycin Resistant *Staphylococcus aureus* infections complex [17].

Of all the enterobacteriaceae isolates, 21.33% (95% CI= 12.19 - 34.64) were identified as Extended spectrum β -lactamase (ESBL) producing pathogens (Table 3). These findings are comparable with several other studies that have reported ESBL prevalence rates in a range of 4 to 46% [10,25]. Co-resistances against therapeutic options for ESBL enterobacteriaceae were reported in 25% (4/16) of all ESBL producers. This translates to the possibility that there is 5% chance that an enterobacteriaceae picked from ward items and surfaces would be an ESBL co-resistant pathogen, nevertheless this slight chance still indicates the existence of ESBL producers in this hospital that would surely present with quite a great patient management problem especially unresponsiveness to what could be used as ESBL treatment alternatives [25,26,28,69].

The occurrence of co-resistance patterns among ESBL producers in the present study is quite consistent with numerous earlier findings that have documented similar co-resistances to non- β -lactam antibiotics [25,36]. Their presence could be attributed to chromosomal mutations conferring resistance to β -Lactam antibiotics and large plasmids that mediate multiple resistance to other antimicrobial classes [67,70,71], that would be used as alternatives against ESBL producers [36,72-74]. In our study, carbapenems

were the most effective antibiotics against enterobacteriaceae including all ESBL and AmpC producers (Table 2), registering a susceptibility rate of 100%, a finding that highlights their importance in the management of ESBL and AmpC producing enterobacteriaceae at Kiwoko Hospital. This is in agreement with several other studies that have recently demonstrated that Carbapenems remain the preferred treatment option against such resistant enterobacteriaceae [28,29,75]. All ESBL producers were sensitive (100%) to Cefoxitin, a common cephamycin. This accords with earlier recommendations of its use as an alternative antimicrobial agent against β -Lactamase producing enterobacteriaceae [28,73].

Out of all the 167 isolates, 25 (16.17%) were detected as Multi-drug resistant species (Table 9). The presence of MDRs on hospital items possibly suggests pathogen shedding from patients who frequently misuse and overuse antibiotics [76,77]. In many countries antibiotics are available over counters without prescription or supervision [78-81]. Similarly, episodes of antimicrobial self-medication among Ugandans have also been reported over the years [82,83]. Such practices would subsequently increase patient management costs and prolonged hospital stays as observed by several earlier scientists [3,5,6,84]. Our findings on MDR pathogens therefore support numerous studies that have cited their alarmingly growing occurrence in Sub-Saharan Africa [85-89].

4. CONCLUSION

In conclusion, findings from our study have revealed the occurrence of resistant bacterial strains on hospital fomites for the first time in our region. In terms of diagnostic accuracy, it is therefore important that routine detection of resistant patterns in the form of MLS_B, ESBL, AmpC, MDR and MRSA from clinical samples is done together with the recommended WHO nosocomial infection control guidelines, to aid in the improvement of patient management strategies, and as a lift towards fighting increasing antimicrobial resistances here in Africa.

It remains unclear as to which extent bacterial contamination of ward surfaces/items contribute to the transmission of pathogens to hospitalized patients. However, evidence based on this study does indicate an urgent need to alert and

educate hospital staff about the potential health risks associated with the use of fomites.

ETHICAL APPROVAL

The study was approved by the research and ethics review committee of Makerere University, school of Biotechnology and Laboratory Science College of Veterinary Medicine, Animal resources and Biosecurity. Permission was sought from the hospital and laboratory authorities. The ethical principles of a scientific research, related laws and regulations were strictly adhered to as well.

ACKNOWLEDGEMENTS

We would like to acknowledge the unfailing support from Ms. Evelyn Ayot, Dr. Asuman Sengooba, Dr. Rose Clarke Nanyonga, and Dr. James Nyonyintono towards this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Garner J, Favero M. Centre for disease prevention and control guidelines for hand-washing and hospital environmental control. *Infection Control*. 1986;7:231-43.
2. Dancer SJ. The role of environmental cleaning in the control of hospital-acquired infection. *The Journal of Hospital Infection*. 2009;73(4):378-85. DOI: 10.1016/j.jhin.2009.03.030 PubMed PMID: 19726106
3. Coello R, Glenister H, Fereres J. The cost of infection in surgical patients: A case-control study. *The Journal of Hospital Infection*. 1993;25:239-50.
4. Allegranzi B, Bagheri Nejad S, Combescure C, Graafmans W, Attar H, Donaldson L, et al. Burden of endemic health-care associated infection in developing countries: Systematic review and meta-analysis. *Lancet*. 2011; 377(9761):228-41. DOI: 10.2471/BLT.11.088179 PubMed PMID: 22084514 PubMed Central PMCID: PMC3209981
5. Kramer A, Schwebke I, Kampf G. How long do nosocomial pathogens persist on inanimate surfaces? A systematic review.

- BMC Infectious Diseases. 2006;6:130-30.
DOI: 10.1186/1471-2334-6-130
PubMed PMID: PMC1564025
6. Madani N, Rosenthal VD, Dendane T, Abidi K, Zeggwagh AA, Abouqal R. Health-care associated infections rates, length of stay and bacterial resistance in an intensive care unit of Morocco: Findings of the International Nosocomial Infection Control Consortium (INICC). *International Archives of Medicine*. 2009;2:29-29.
DOI: 10.1186/1755-7682-2-29
PubMed PMID: PMC2765432
 7. Ducei G, Fabry J, Nicolle L, editors. *Prevention of hospital acquired infections, a practical guide*. 2nd ed. Geneva Switzerland: World Health Organization; 2002.
 8. Mayon-White R. An international survey of the prevalence of hospital-acquired infection. *J Hosp Infect*. 1988;11:43-48.
 9. World Health Organization. *Guidelines on hand hygiene in health care (advanced draft): A summary*. Geneva: World Health Organisation; 2005.
 10. Rothe C, Schlaich S, Thompson S. Healthcare-associated infections in sub-Saharan Africa. *Journal of Hospital Infection*. 2013;85:257-67.
DOI: 10.1016/j.jhin.2013.09.008
 11. Jroundi I, Khoudri I, Azzouzi A. Prevalence of hospital acquired infection in a Moroccan hospital. *American Journal of Infection Control*. 2007;(35):412-16.
 12. Trillis F, Eckstein EC, Budavich R, Pultz MJ, Donskey CJ. Contamination of hospital curtains with healthcare-associated pathogens. *Infection Control and Hospital Epidemiology*. 2008;29(11):1074-6.
DOI: 10.1086/591863
PubMed PMID: 18823274
 13. Catano JC, Echeverri LM, Szela C. Bacterial contamination of clothes and environmental items in a third-level hospital in Colombia. *Interdisciplinary Perspectives on Infectious Diseases*. 2012;2012:507-640.
DOI: 10.1155/2012/507640
PubMed Central PMCID: PMC3321286
 14. Hayden MK, Blom DW, Lyle EA, Moore CG, Weinstein RA. Risk of hand or glove contamination after contact with patients colonized with vancomycin-resistant *Enterococcus* or the colonized patients' environment. *Infection Control and Hospital Epidemiology*. 2008;29(2):149-54.
DOI: 10.1086/524331
PubMed PMID: 18179370
 15. Feglo P, Afriyie-Asante A. Environmental impact on postoperative wound infections in a privately owned hospital in Ghana. *African Journal of Microbiology Research*. 2014;8(15):1620-26.
DOI: 10.5897/ajmr2013.6438
 16. Boyce JM. Environmental contamination makes an important contribution to hospital infection. *The Journal of Hospital Infection*. 2007;65(Suppl 2):50-4.
DOI: 10.1016/S0195-6701(07)60015-2
PubMed PMID: 17540242
 17. Drees M, Snyderman DR, Schmid CH, Barefoot L, Hansjosten K, Vue PM, et al. Prior environmental contamination increases the risk of acquisition of vancomycin-resistant enterococci. *Clinical infectious diseases: An official publication of the Infectious Diseases Society of America*. 2008;46(5):678-85.
DOI: 10.1086/527394
PubMed PMID: 18230044
 18. Maki DG, Alvarado CJ, Hassemer CA, Zilz MA. Relation of the inanimate hospital environment to endemic nosocomial infection. *The New England Journal of Medicine*. 1982;307(25):1562-6.
DOI: 10.1056/NEJM198212163072507
PubMed PMID: 6815529
 19. Williams C, Davis DL. Methicillin-resistant *Staphylococcus aureus* fomite survival. *Clinical laboratory science. Journal of the American Society for Medical Technology*. 2009;22(1):34-8.
PubMed PMID: 19354027
 20. Sasahara T, Hayashi S, Morisawa Y, Sakihama T, Yoshimura A, Hirai Y. *Bacillus cereus* bacteremia outbreak due to contaminated hospital linens. *European Journal of Clinical Microbiology & Infectious Diseases*. 2011;30(2):219-26.
DOI: 10.1007/s10096-010-1072-2
 21. Wiener-Well Y, Galuty M, Rudensky B, Schlesinger Y, Attias D, Yinnon AM. Nursing and physician attire as possible source of nosocomial infections. *American Journal of Infection Control*. 2011;39(7):555-9.
DOI: 10.1016/j.ajic.2010.12.016
PubMed PMID: 21864762
 22. Sing-on T, Wen-sen L, Tsong-Yih O, Yu-Chia H, Wuan-Chan L, Yi-Chun L. Bacterial contamination of patients' medical charts in a surgical ward and the intensive care unit: Impact on nosocomial

- infections, Wan Fang Hospital. J Microbiol Immunol Infect. 2009;42(1):86-91.
23. Bhalla A, Pultz NJ, Gries DM. Acquisition of nosocomial pathogens on hands after contact with environmental surfaces near hospitalized patients. Infection Control and Hospital Epidemiology. 2004;25(2):164-67.
 24. French GL, Otter JA, Shannon KP, Adams NM, Watling D, Parks MJ. Tackling contamination of the hospital environment by methicillin-resistant *Staphylococcus aureus* (MRSA): A comparison between conventional terminal cleaning and hydrogen peroxide vapour decontamination. J Hosp Infect. 2004;57:31-37.
 25. Kiratisin P, Chattammanat S, Sa-Nguansai S, Dansubutra B, Patthamalai P, Nangpatharapornthawee P, et al. A 2-year trend of extended-spectrum - lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* in Thailand: An alert for infection control. Royal Society of Tropical Medicine and Hygiene. 2008;102:460-64. DOI: 10.1016/j.trstmh.2008.02.005
 26. Grover N, Sahni AK, Bhattacharya S. Therapeutic challenges of ESBLs and AmpC beta-lactamase producers in a tertiary care center. Medical Journal Armed Forces India. 2013;69(1):4-10. DOI: 10.1016/j.mjafi.2012.02.001 PubMed PMID: PMC3862620
 27. Madhumati B, Rani L, Ranjini CY, Rajendran R. Prevalence of AMPC beta lactamases among gram negative bacterial isolates in a tertiary care hospitals. Int J Curr Microbiol App Sci. 2015;4(9):219-27.
 28. Kanj SS, Kanafani ZA. Current concepts in antimicrobial therapy against resistant gram-negative organisms: Extended-spectrum β -lactamase-producing enterobacteriaceae, carbapenem-resistant enterobacteriaceae, and multidrug-resistant *Pseudomonas aeruginosa*. Mayo Clinic Proceedings. 2011;86(3):250-59. DOI: 10.4065/mcp.2010.0674 PubMed PMID: PMC3046948
 29. Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: An international expert proposal for interim standard definitions for acquired resistance. Clinical Microbiology and Infection. 2012;18(3):268-81. DOI: 10.1111/j.1469-0691.2011.03570.x
 30. Mittal R, Aggarwal S, Sharma S, Chhibber S, Harjai K. Urinary tract infections caused by *Pseudomonas aeruginosa*: A minireview. Journal of Infection and Public Health. 2009;2:101-11. DOI: 10.1016/j.jiph.2009.08.003
 31. Zmantar T, Kouidhi B, Miladi H, Bakhrouf A. Detection of macrolide and disinfectant resistance genes in clinical *Staphylococcus aureus* and coagulase-negative staphylococci. BMC Research Notes. 2011;4:453-53. DOI: 10.1186/1756-0500-4-453 PubMed PMID: PMC3212975
 32. Yilmaz G, Aydin K, Iskender S, Caylan R, Koksali I. Detection and prevalence of inducible clindamycin resistance in staphylococci. Journal of Medical Microbiology. 2007;56(3):342-45. DOI: 10.1099/jmm.0.46761-0
 33. Lall M, Sahni AK. Prevalence of inducible clindamycin resistance in *Staphylococcus aureus* isolated from clinical samples. Medical Journal Armed Forces India. 2014;70:43-47. DOI: org/10.1016/j.mjafi.2013.01.004
 34. Bauer AW, Kirby WM, Sherris JC, Turck H. Antibiotic susceptibility testing by a standard single disk method. Am J clin Pathol. 1966;45(4):493-96.
 35. Clinical and Laboratory Standards Institute. Performance Standards for antimicrobial susceptibility testing; Twenty-Fourth Informational Supplement. CLSI document M100-S24. ISBN 1-56238-897-5 [Print] ISBN 1-56238-898-3 [Electronic]. 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087 USA: Clinical and Laboratory Standards Institute; 2014.
 36. Paterson DL, Bonomo RA. Extended-spectrum beta-lactamases: A clinical update. Clinical Microbiology Reviews. 2005;18(4):657-86. DOI: 10.1128/cmr.18.4.657-686.2005
 37. Parveen MR, Harish BN, Parija SC. AmpC beta lactamases among Gram negative clinical isolates from a tertiary hospital, South India. Brazilian Journal of Microbiology. 2010;41:596-602.
 38. Burgess DS, Nathisuwan S. Cefepime, piperacillin/tazobactam, gentamicin, ciprofloxacin, and levofloxacin alone and in combination against *Pseudomonas aeruginosa*. Diagnostic Microbiology and Infectious Disease. 2002;44(1):35-41.

39. Fiebelkorn KR, Crawford SA, McElmeel ML, Jorgensen JH. Practical disk diffusion method for detection of inducible clindamycin resistance in *Staphylococcus aureus* and coagulase-negative staphylococci. *Journal of Clinical Microbiology*. 2003;41(10):4740-44. DOI: 10.1128/JCM.41.10.4740-4744.2003 PubMed PMID: PMC254362
40. Nwankwo E. Isolation of pathogenic bacteria from fomites in the operating rooms of Murtala Mohammed Specialist Hospital, Kano. *Pan African Medical Journal*. 2012;1:12-90.
41. Ensayef S, Al-Shalchi S, Sabbar M. Microbial contamination in the operating theatre: A study in a hospital in Baghdad. *East Mediterr Health Journal*. 2009;15(1):219-23.
42. Gelaw A, Gebre-Selassie S, Tiruneh M, Mathios E, Yifru S. Isolation of bacterial pathogens from patients with postoperative surgical site infections and possible sources of infections at the University of Gondar Hospital, Northwest Ethiopia. *J Environ Occup Sci*. 2014;3(2):103-08.
43. Hold A. Infection control in theatre. *South Afr J Anaesth Analg*. 2011;17(1):56-64.
44. Spagnolo AM, Ottria G, Amicizia D, Perdelli F, Cristina ML. Operating theatre quality and prevention of surgical site infections. *Journal of Preventive Medicine and Hygiene*. 2013;54(3):131-37. PubMed PMID: PMC4718372
45. Aswin RS, Hegde A. Bacterial contamination of stethoscopes. *International Journal of Scientific Research*. 2014;3(11):406-08. DOI: 10.15373/22778179
46. Qaday J, Sarike M, Mwakyoma A, Kifaro E, Mosha D, Tarimo R. Bacterial contamination of Medical doctors and students' white coats at Kilimanjaro Christian Medical Centre, Moshi, Tanzania. *International Journal of Bacteriology*. 2015;2015. DOI: 10.1155/2015/507890
47. Gebremariam T, Declaro M. Operating theaters as a source of nosocomial infection: A systematic review. 2014;3(1):5-8. DOI: 10.4103/2278-0521.130196
48. Greco D, Magombe I. Hospital acquired infections in Lacor Hospital Uganda. *J Prev Med Hyg*. 2011;52:55-58.
49. Centers for Disease Control and Prevention NaSH-AIPR. National and State healthcare associated infections: progress report; 2016. Available:<http://www.cdc.gov/hai/progress-report/index.html>
50. De Giallully C, Morange V, De Giallully E, Loulergue J, Van der Mee N, Quentin R. Blood pressure cuff as potential vector of pathogenic microorganisms: A prospective study in a teaching hospital. *Infection Control and Hospital Epidemiology*. 2006;27(9):940-43.
51. Riggs MM, Sethi AK, Zabarsky TF, Eckstein EC, Jump RL, Donskey CJ. Asymptomatic carriers are a potential source for transmission of epidemic and nonepidemic clostridium difficile strains among long-term care facility residents. *Clinical Infectious Diseases*. 2007; 45(8):992-98. DOI: 10.1086/521854 PubMed PMID: 17879913
52. Maryam A, Hadiza US, Aminu UM. Characterization and determination of antibiotic susceptibility pattern of bacteria isolated from some fomites in a teaching hospital in Northern Nigeria. *African Journal of Microbiology Research*. 2014;8(8):814-18. DOI: 10.5897/AJMR2013.6512
53. Vandepitte J, Verhaegen J, Engbaek K, Rohner P, Piot P, Heuck CC. *Basic Laboratory Procedures in Clinical Bacteriology*. 2nd ed. Geneva, Switzerland: World Health Organisation; 2003.
54. Cogen AL, Nizet V, Gallo RL. Skin microbiota: A source of disease or defence? *The British Journal of Dermatology*. 2008;158(3):442-55. DOI: 10.1111/j.1365-2133.2008.08437.x PubMed PMID: PMC2746716
55. Cheesbrough M. *District laboratory practice in tropical countries*. 2 ed. Newyork: Cambridge University Press; 2006.
56. Iwatsuki K, Yamasaki O, Morizane S, Oono T. Staphylococcal cutaneous infections: Invasion, evasion and aggression. *Journal of Dermatological Science*. 2006;42(3):203-14. DOI: 10.1016/j.jdermsci.2006.03.011 PubMed PMID: 16679003
57. Sakwinska O, Blanc DS, Lazor-Blanchet C, Moreillon M, Giddey M, Moreillon P. Ecological temporal stability of *Staphylococcus aureus* nasal carriage.

- Journal of Clinical Microbiology. 2010;48(8):2724-28.
DOI: 10.1128/JCM.02091-09
PubMed PMID: PMC2916613
58. Al-Zoubi MS, Al-Tayyar IA, Hussein E, Jabali AA, Khudairat S. Antimicrobial susceptibility pattern of *Staphylococcus aureus* isolated from clinical specimens in Northern area of Jordan. Iranian Journal of Microbiology. 2015;7(5):265-72.
PubMed PMID: PMC4695508
 59. Peacock SJ, de Silva L, Lowy FD. What determines nasal carriage of *Staphylococcus aureus*? Trends Microbiol. 2001;9:605-10.
 60. Fraimow HS. Systemic antimicrobial therapy in osteomyelitis. Seminars in Plastic Surgery. 2009;23(2):90-99.
DOI: 10.1055/s-0029-1214161
PubMed PMID: PMC2884905
 61. Elyasi S, Khalili H, Dashti-Khavidaki S, Mohammadpour A. Vancomycin-induced nephrotoxicity: Mechanism, incidence, risk factors and special populations. A literature review. European Journal of Clinical Pharmacology. 2012;68(9):1243-55.
DOI: 10.1007/s00228-012-1259-9
PubMed PMID: 22411630.
 62. van Hal SJ, Paterson DL, Lodise TP. Systematic review and meta-analysis of vancomycin-induced nephrotoxicity associated with dosing schedules that maintain troughs between 15 and 20 milligrams per liter. Antimicrobial Agents and Chemotherapy. 2013;57(2):734-44.
DOI: 10.1128/aac.01568-12
 63. Liu C, Bayer A, Cosgrove SE, Daum RS, Fridkin SK, Gorwitz RJ, et al. Clinical practice guidelines by the infectious diseases society of america for the treatment of methicillin-resistant *Staphylococcus aureus* infections in adults and children. Clinical infectious diseases: an official publication of the Infectious Diseases Society of America. 2011;52(3):18-55.
DOI: 10.1093/cid/ciq146
PubMed PMID: 21208910
 64. Siberry GK, Tekle T, Carroll K, Dick J. Failure of clindamycin treatment of methicillin resistant resistant *Staphylococcus aureus* expressing inducible clindamycin resistance *in vitro*. Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America. 2003;37:1257-60.
 65. Tsuji BT, Rybak MJ, Cheung CM, Amjad M, Kaatz GW. Community- and health care-associated methicillin-resistant *Staphylococcus aureus*: A comparison of molecular epidemiology and antimicrobial activities of various agents. Diagnostic Microbiology and Infectious Disease. 2006;58:41-47.
DOI: 10.1016/j.diagmicrobio.2006.10.021
 66. Hiramatsu K. Vancomycin-resistant *Staphylococcus aureus*: A new model of antibiotic resistance. Lancet Infectious Diseases. 2001;1:147-55.
 67. Alekshun MN, Levy SB. Molecular mechanisms of antibacterial multidrug resistance. Cell. 2007;128(6):1037-50.
DOI: 10.1016/j.cell.2007.03.004
 68. De Lencastre H, Oliveira D, Tomasz A. Antibiotic resistant *Staphylococcus aureus*: A paradigm of adaptive power. Current Opinion in Microbiology. 2007;10:428-35.
DOI: 10.1016/j.mib.2007.08.003
 69. Nasir KM, Preeti S, Vikili C, Singh NP. Prevalence of ESBL and AmpC β -lactamase in gram negative bacilli in various clinical samples at tertiary care hospital. Int Res J Medical Sci. 2015;3(8):1-6.
 70. Canton R, Novais A, Valverde A, Machado E, Peixe L, Baquero F, et al. Prevalence and spread of extended-spectrum β -lactamase-producing enterobacteriaceae in Europe. Clinical microbiology and infection: The official publication of the European Society of Clinical Microbiology and Infectious Diseases. 2008;14(Suppl. 1):144-53.
 71. Schwaber MJ, Navon-Venezia S, Schwartz D, Carmeli Y. High levels of antimicrobial co-resistance among extended-spectrum- β -lactamase-producing enterobacteriaceae. Antimicrobial Agents and Chemotherapy. 2005;49(5):2137-39.
DOI: 10.1128/AAC.49.5.2137-2139.2005
 72. van Hoek AHAM, Mevius D, Guerra B, Mullany P, Roberts AP, Aarts HJM. Acquired antibiotic resistance genes: An overview. Frontiers in Microbiology. 2011;2:203.
DOI: 10.3389/fmicb.2011.00203
 73. Rakotonirina HC, Garin B, Randrianirina F, Richard V, Talarmin A, Arlet G. Molecular characterization of multidrug-resistant extended-spectrum β -lactamase-producing enterobacteriaceae isolated in

- Antananarivo, Madagascar. *BioMed Central Microbiology*. 2013;13(85). DOI: 10.1186/1471-2180-13-85
74. Nikaido H. Multidrug resistance in bacteria. *Annual Review of Biochemistry*. 2009;78:119-46. DOI:10.1146/annurev.biochem.78.082907.145923. PubMed PMID: PMC2839888
 75. Chelliah A, Ravinder T, Katragadda R, Leela KV, Babu RN. Isolation of MRSA, ESBL and AmpC - beta -lactamases from neonatal sepsis at a tertiary care hospital. *Journal of clinical and diagnostic research*. 2014;8(6):24-27. DOI: 10.7860/jcdr/2014/8597.4512
 76. Cars O, Nordberg P. Antibiotic resistance-the faceless threat. *International Journal of Risk and Safety in Medicine*. 2005;17:103-10.
 77. Davies J, Davies D. Origins and evolution of antibiotic resistance. *Microbiology and Molecular Biology Reviews*. 2010;74(3):417-33. DOI: 10.1128/MMBR.00016-10
 78. Richman P, Garra G, Eskin B, Nashed A, Cody R. Oral antibiotic use without consulting a physician: A survey of ED patients. *Am J Emerg Med*. 2001;19. DOI: 10.1053/ajem.2001.20035
 79. Skliros E, Merkouris P, Papazafiropoulou A, Gikas A, Matzouranis G, Papafragos C, et al. Self-medication with antibiotics in rural population in Greece: A cross-sectional multicenter study. *BMC Family Practice*. 2010;11(1):1-3. DOI: 10.1186/1471-2296-11-58
 80. Aqeel T, Shabbir A, Basharat H, Bukhari M, Mobin S, Shahid H, et al. Prevalence of self-medication among urban and rural population of Islamabad, Pakistan. *Tropical Journal of Pharmaceutical Research*. 2014;13(4):627-33.
 81. Borg MA, Scicluna EA. Over-the-counter acquisition of antimicrobial drugs in the maltese general population. *Inter J Antimicrob Agents*. 2002;20. DOI: 10.1016/s0924-8579(02)00194-2
 82. Ocan M, Bwanga F, Bbosa GS, Bagenda D, Waako P, Okeng JO, et al. Patterns and Predictors of Self-Medication in Northern Uganda. *PLoS ONE*. 2014;9(3). DOI: 10.1371/journal.pone.0092323
 83. Anyama N, Adome RO. Community pharmaceutical care: An 8-month critical review of two pharmacies in Kampala. *African Health Sciences*. 2003;3(2):87-93. PubMed PMID: PMC2141595
 84. Chen YY, Chou YC, Chou P. Impact of nosocomial infection on cost of illness and length of stay in intensive care units. *Infect Control Hosp Epidemiol*. 2005;26:281-87.
 85. Leopold SJ, Van Leth F, Tarekegn H, Schultsz C. Antimicrobial drug resistance among clinically relevant bacterial isolates in sub-Saharan Africa: A systematic review. *Journal of Antimicrobial Chemotherapy*; 2014. DOI: 10.1093/jac/dku176
 86. Ibrahim ME, Bilal NE, Hamid ME. Increased multi-drug resistant *Escherichia coli* from hospitals in Khartoum state, Sudan. *African Health Sciences*. 2012;12(3):368-75. PubMed PMID: PMC3557680
 87. Okeke IN, Aboderin OA, Byarugaba DK, Ojo KK, Opintan JA. Growing problem of multidrug-resistant enteric pathogens in Africa. *Emerging Infectious Diseases*. 2007;13(11):1640-46. DOI: 10.3201/eid1311.070674 PubMed PMID: PMC3375797
 88. Godebo G, Kibru G, Tassew H. Multidrug-resistant bacterial isolates in infected wounds at Jimma University Specialized Hospital, Ethiopia. *Annals of Clinical Microbiology and Antimicrobials*. 2013;12:17-17. DOI: 10.1186/1476-0711-12-17 PubMed PMID: PMC3724577
 89. Kimang'a AN. A situational analysis of antimicrobial drug resistance in africa: Are we losing the battle? *Ethiopian Journal of Health Sciences*. 2012;22(2):135-43. PubMed PMID: PMC3407837

© 2016 Segujja et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
 The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/10035>